

Abstract No. Flin0615

Relative Metal Concentrations in Amyloid Plaques of Alzheimer's Disease, as Determined by Synchrotron X-ray Fluorescence

J. M. Flinn, D.H. Linkous (George Mason U.), A.L. Friedlich (Mass. General), A. Lanzirotti (U. of Chicago), W. Rao (NSLS), K. Newell (Mass General), C.J. Frederickson (NeuroBioTex), B.F. Jones (USGS), P. Bertsch (Savannah River Ecology Lab), & A.I. Bush (Mass General)

Beamline(s): X26A

Introduction: Alzheimer's disease amyloid plaques have previously been identified as containing higher metal concentrations than surrounding tissue [1]. However, the distribution of metals within plaques has not been mapped. Synchrotron X-Ray fluorescence allows the measurement of metal distributions within biological tissue. Preliminary analyses of metal concentrations and distribution within plaques of different morphologies are reported here.

Methods and Materials: Post-mortem human brain slices were obtained from stage 6 Alzheimer's disease patients. Tissue was fresh, unfixed, and sectioned to 20 μ m thickness. X-ray synchrotron fluorescence was used to examine the distribution of metals within the tissue. Potassium, iron, copper and zinc levels were examined in different types of amyloid deposits: a congophilic vascular amyloid region and two diffuse plaques. Copper and zinc levels were also measured in a dense core plaque; levels for all metals were also measured in non-plaque tissue.


Results: Iron levels were ~ 3 x background in all cases, although the concentration in the center of the diffuse core plaque (3.9 x background) was greater than in the periphery (3.1 x background). Both zinc and potassium levels were ~ 5 x higher than background in the center of the vascular amyloid region, and zinc levels were also ~ 5 x background in the center of the dense core plaque (see Fig 1). Potassium and zinc levels in the periphery of the region were 1.9 and 1.6 x background, respectively. Copper levels were 1.3 x background in all regions of the vascular and diffuse plaques, but were ~ 4 x background in the center of the dense core plaque (see Fig 1).

Conclusions: These preliminary data indicate that metals vary not only within plaques, but also across plaque types.

Acknowledgments: This study was supported by the Harvard Brain Resource Center (PHS MH/NS 31862); a BNL general user grant for NSLS beamline access (J.M.F.); a George Mason University Research Assistantship (D.H.L.); F32NS10874 (A.L.F.); NS40215, NS46682 (C.J.F.); 2R01AG12686, and an Alzheimer's Association IIRG (A.I.B.). National Synchrotron Light Source (NSLS) is supported by the U.S. Department of Energy under Contract No. DE-AC02-76CH00016.

References:

[1]. M.A. Lovell, J.D. Robertson, W.J. Teesdale, J.L. Campbell, and W.R. Markesbery, "Copper, iron and zinc in Alzheimer's disease senile plaques," *J. Neurol. Science*, **158** (1), 47-52 (1998).



Dense Core Plaque: Cu

Dense Core Plaque: Zn

Figure 1